

ENHANCED OUTCROP TO THE SUBSURFACE CORRELATION IN THE VACA MUERTA FORMATION

Taury Smith¹, Ralf J. Weger, and Gregor P. Eberli
¹⁾ Smith Stratigraphic LLC

PROJECT OBJECTIVES

- Tie the outcrop to the subsurface based on the composition in thin sections and cuttings from 20 wells and outcrop sections across the basin.
- Evaluate the similarities and differences of compositional variations in the sequences and cycles to capture the lateral and vertical facies variability within the Vaca-Muerta Formation.
- Provide a (semi)quantitative comparison of facies similarities in the outcrop and subsurface.

PROJECT RATIONALE

In the past we have successfully correlated the outcrop successions of the Vaca Muerta Formation in the Neuquén Basin to the subsurface with geophysical and geochemical data. Synthetic seismic sections constructed from facies and geometries observed in the Picun Leufu area and the Sierra de




Figure 1: Correlation of cycles and sequences between three wells using cuttings and cores together with log information. The quantification of the composition adds valuable information about the lateral and vertical changes of facies.

la Vaca Muerta (SdIVM) are very similar to the geometries observed on seismic sections (Zeller et al., 2015 a, b). The correlation of outcrop sections to core and well logs is achieved by adding spectral gamma ray, TOC and other geochemical properties with high resolution (0.5-1 m spacing) to the lithology in each section. Key log-signature subdivisions from the subsurface correspond to sequence stratigraphic divisions identified in the outcrop sections (Eberli et al., 2017).

With these correlations in hand an enhanced correlation is possible that takes into account facies variations within the cycles and sequences. This correlation is based on a quantitative analysis of thin sections from outcrop successions and its comparison to the subsurface data (Fig. 1). The advantage of this facies based correlation is that it captures the sedimentological variations that are created by the cyclic deposition. In addition, it reflects the proximal distal facies changes as well as changes through time.

APPROACH

The study can rely on a large data set from the subsurface produced by Taury Smith which will be compared to the outcrop sections. The data set consists of a cutting and core study of more than 20 wells across the basin from the top of the Quintuco to the base of the Vaca Muerta using more than 5000 thin sections. Based on the cuttings and logs detailed stratigraphic columns are constructed that allow sequences and cycles to be picked (Fig. 1). Taury Smith will work with us to assemble a similar data set in our outcrop sections so that they can be compared to the subsurface. Some wells he studied are within 10 km of the outcrop belt.

SIGNIFICANCE

This facies-based correlation of outcrop to subsurface will add valuable information regarding the facies changes between the outcrop at the western side of the basin and subsurface areas further to the east. If the comparison reveals strong similarities, it will strengthen the potential of the outcrops to act as analogs for the subsurface. Variations would indicate lateral facies changes along the clinofolds in the basin.

REFERENCES

- Eberli, G.P., Weger, R.J., Tenaglia, M., Rueda, L., Rodriguez Blanco, L., Zeller, M., McNeill, D.F., Murray, S., and Swart, P.K., 2017, The Unconventional Play in the Neuquén Basin, Argentina – Insights from the Outcrop for the Subsurface. Extended Abstract, URTEC 2687581 July 24-26, 2017 Austin, TX.
- Zeller, M., K. Reid, S.B., Eberli, G.P., Weger, R.J., and Massaferro, J.L., 2015a. Sequence architecture and heterogeneities of a field Scale Vaca Muerta analog (Neuquén Basin, Argentina) - From outcrop to synthetic seismic. *Marine and Petroleum Geology*, v. 66, p.829-847.
- Zeller, M., Verwer K., Eberli, G.P., Massaferro, J.L., Schwarz, E., and Spalletti, L., 2015b. Depositional controls on mixed carbonate-siliciclastic cycles and sequences on gently inclined shelf profiles: *Sedimentology*, v. 62, p. 2009-2037.